

RESEARCH REPORT

In vitro hemotoxic, α -neurotoxic and vasculotoxic effects of the Mexican black-tailed rattlesnake (*Crotalus molossus nigrescens*) venom

David Meléndez-Martínez¹, Eduardo Macías-Rodríguez¹, Rodrigo Vázquez-Briones¹, Estuardo López-Vera², Martha Sandra Cruz-Pérez³, Alejandra Vargas-Caraveo¹, Ana Gatica-Colima¹, Luis Fernando Plenge-Tellechea^{1,*}

¹Laboratorio de Biología Molecular y Bioquímica, Departamento de Ciencias Químico Biológicas, Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, Chihuahua, México, ²Laboratorio de Toxinología Marina, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Circuito Exterior s/n, Ciudad Universitaria, 04510 Coyoacán, Ciudad de México, México, ³Herpetario de la Universidad Autónoma de Querétaro, Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Juriquilla, Querétaro, México

*Correspondence to: Luis Fernando Plenge-Tellechea, Email: fplenge@uacj.mx, Tel/Fax: +52 656 6881800 ext 1621

Received: 02 December 2016; Revised: 03 March 2017; Accepted: 09 March 2017; Published: 29 March 2017

© Copyright The Author(s). This is an open access article, published under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>). This license permits non-commercial use, distribution and reproduction of this article, provided the original work is appropriately acknowledged, with correct citation details.

ABSTRACT

The Mexican black-tailed rattlesnake *Crotalus molossus nigrescens* is distributed in the Mexican plateau. Its venom is known to cause hemolysis and presents fibrinogen coagulase, collagenase and fibrinolytic activities. These activities may be associated with hemostatic alterations, such as platelet aggregation, hemolysis and fibrinolysis, often described in ophidic accidents. However, the mechanisms of action of the *C. m. nigrescens* venom remain unclear. In this study we investigated the *in vitro* hemotoxic, neurotoxic, and vasculotoxic effects of the venom. We found that this venom produces two types of hemolytic responses, Oxyhemoglobin release and Methemoglobin formation. As a result of the cytotoxicity to endothelial cells produces morphological biphasic toxicity. The first step in this process is characterized by morphological changes, as well as the loss of cellular adhesion and reduction in thickness. The second phase is characterized by massive cellular aggregation and death. It also induced laminin, type IV collagen, perlecan and nidogen degradation. However, the venom did not modulate the muscular fetal and neuronal nicotinic acetylcholine receptors activity. Thus, we concluded that the *C. m. nigrescens* venom produced hemolysis and hemorrhages via degradation of the basement membrane components and endothelial cell cytotoxicity, but not by neurotoxicity at the receptor level in nicotinic acetylcholine receptors.

KEYWORDS: *Crotalus molossus nigrescens*, hemolysis, hemorrhage, neurotoxicity, snake venom

INTRODUCTION

The Mexican black-tailed rattlesnake *Crotalus molossus nigrescens* is found in the Mexican plateau from Southern Chihuahua to North of Oaxaca (Lemos-Espinal and Smith, 2009), and is morphologically characterized by a black coloration in the tail. As in other venomous snakes, its venom is a complex mix of proteins, such as metalloproteinases (SVMP), serine proteinases, phospholipases A2 (PLA2), L-aminoacid oxidases (LAAO), disintegrins, C-type lectins, Cys-rich secretory proteins and other minor

proteases toxins. This suggests that the toxicity of this venom is of predominantly proteolytic nature, such as fibrinogenolysis, fibrinolysis, hemorrhages and platelet aggregation (Mackessy, 2010). *Crotalus molossus* venom generates hemostatic alterations as platelet aggregation, hemolysis and fibrinolysis, as well as hemorrhages and necrosis (Hardy et al, 1982; Yarema and Curry, 2005).

The Mexican black-tailed rattlesnake venom is known to have protease, phospholipase, phosphodiesterase, deoxyribonuclease, fibrinogen coagulase, collagenase and

fibrinolytic activities; only two toxins have been isolated, proteinase E, a 21.39kDa SVMP and 75kDa thrombin-like serineproteinase (Ramírez et al, 1990). *In vitro* studies demonstrate that *C. m. nigrescens* venom is more hemolytic than *Crotalus atrox* and *Crotalus tigris* venoms (Macias-Rodríguez et al, 2014) via plasmatic membrane degradation catalyzed by PLA2. The fibrinogen and collagen degradation previously mentioned suggest that *C. m. nigrescens* venom could be involved in hemorrhage formation and hemostatic alterations (Gutiérrez et al, 2005). Neurotoxicity in *Crotalus* genera has been described for two PLA2 toxins: i) Crotoxin, isolated from *Crotalus durissus* venom that can function as a presynaptic neurotoxin, inhibiting the chlorine uptake in neurons (Kattah et al, 2000), and a postsynaptic neurotoxin, modulating the nicotinic acetylcholine receptor (Bon et al, 1979); and ii) Mojave toxin, isolated from *Crotalus scutulatus* venom, can act as a Calcium channel inhibitor (Valdes et al, 1989) but there is no clear evidence of neurotoxicity in *C. m. nigrescens* venom. Thus due to a lack of data on *C. m. nigrescens* venom toxicity, this study was designed to investigate venom hemotoxicity, α -neurotoxicity and vasculotoxicity using *in vitro* techniques.

MATERIAL AND METHODS

Venom

Venom samples were obtained from 13 *C. m. nigrescens* specimens maintained at the Universidad Autónoma de Querétaro Herpetary. Venom extraction was performed manually as described previously (Meléndez-Martínez et al, 2014). The *C. m. nigrescens* venom was pooled, lyophilized and stored at -20°C until used. Protein concentration in venom was measured by Lowry protein assay (Lowry et al, 1951) using bovine serum albumin as protein standard.

Hemotoxicity

Blood collection

Blood was collected from at least three healthy, O+ blood type donors, who had not taken any medication in the previous 48hr. Blood samples were collected in BD Vacutainer® sodium heparin tubes for leukocyte cytotoxicity assay and buffered sodium citrate tubes for hemolytic activity assay. Blood collected in heparinized tubes were allowed to separate in the leukocyte rich plasma from Red Blood Cells (RBC) by gravity (Verma and Babu, 1995) and tubes were maintained at room temperature for $\sim 40\text{min}$ at 45 degree inclination.

Hemolytic activity

Hemolytic activity was determined according to Das et al (2013) with some modifications. RBCs were isolated from blood collected in buffered sodium citrate tubes, washed 5 times by centrifugation at 1100xg for 15min, and resuspended in 0.9% (w/v) saline to a final concentration of 10% (v/v) RBC. *Crotalus m. nigrescens* venom concentrations tested (0-640 $\mu\text{g/ml}$) were incubated for 24hr to give sufficient time for hemolysis, at 37°C with 37.5 μl of 10% RBC to a final volume of 500 μl with 0.9% (w/v) saline. Then, the tubes were centrifuged at 6400xg for 10min and supernatant was measured at 540nm for oxyhemoglobin (Oxy-Hb) release and 630nm for methemoglobin (Met-Hb) formation in a Helios Omega UV-Vis spectrophotometer (Thermo Scientific, USA). Distilled H_2O was used as positive control of Oxy-Hb and H_2O_2 as positive control of Met-Hb.

Leukocyte cytotoxicity

Leukocyte rich plasma was washed three times at 500xg for 30min with PBS and resuspended in 5ml of DMEM (high glucose) media supplemented with 10% (v/v) fetal bovine serum and Penicillin-Streptomycin 100U/ml. Leukocyte cultures of 1×10^6 cells were prepared and exposed to




Figure 1. β - and α -hemolytic activity of *C. m. nigrescens* venom. Erythrocytes were treated with different concentrations of *C. m. nigrescens* venom (0-640 $\mu\text{g/ml}$) for 24hr. Data are represented as the mean of three independent experiments with their respective standard error.

different *C. m. nigrescens* venom concentrations (0-10 μ g/ml). The cells were incubated with 5% (v/v) CO₂ at 37°C, at different time intervals (2-12hr). The cell counting was carried out in a Neubauer chamber by diluting the cells 1:1 with 0.1% (w/v) Evan's blue dye.

α -Neurotoxicity

RP-HPLC

Lyophilized *C. m. nigrescens* venom was reconstituted in 0.1% (v/v) trifluoroacetic acid (TFA) and 4.5% (v/v) acetonitrile (ACN) solution. 250 μ g of reconstituted venom was fractionated with a GraceVydac C18 analytical reverse phase column (218TP54, 5 μ m, 4.6x250mm) equipped with a Vydac C18 guard column (218GK54, 5 μ m, 4.6x10mm).

Solution A was 0.1% (v/v) aqueous TFA, and solution B was 0.085% (v/v) TFA in 90% aqueous ACN. Upon injection of the sample, a linear gradient from 5 to 100% solution during 95min was applied at flow rate of 1ml/min. Venom fractions were manually collected and monitored at 280nm, the solvent was eliminated by vacuum concentration, dried fractions were dissolved in water and protein concentration was measured at 280nm by spectrophotometry.

Phospholipase A2 assay

The PLA2 enzymatic activity was performed in egg yolk plates. A 10% (v/v) egg yolk solution was prepared using phospholipase buffer (0.1M Tris-HCl, pH 7.5, 5mM CaCl₂, and 0.5% v/v, Triton X-100), stirring by 10min and




Figure 2. *C. m. nigrescens* venom cytotoxicity on human Leukocytes. **A.** Leukocytes were treated with different concentrations of *C. m. nigrescens* venom (0-10 μ g/ml) at different time intervals (2-12hr). **B.** CC₅₀ (μ g/ml) are shown for every time interval. Data are represented as the mean of three independent experiments with their respective standard error.




Figure 3. Chromatographic fractionation of *C. m. nigrescens* venom by means of an analytical RP C18 HPLC column. Fractions were eluted using a linear gradient of 5-100% solution B (green line) at a flow rate of 1ml/min.

centrifuged at 180xg for 5min to eliminate the non-dissolved egg yolk. The solution was poured into plastic Petri dishes with final concentration of 1% (v/v) of egg yolk solution with 0.5% (w/v) agarose, 0.1M Tris-HCl, 1mM CaCl₂, 0.04% (w/v) rhodamine 6G and 0.5% (v/v) Triton-X100, pH 7.95. *Crotalus m. nigrescens* crude venom and each fraction collected by RP-HPLC were diluted in 5μl of H₂O (~200μg of toxin) and placed into wells in egg yolk plates at 37°C by 24hr. PLA2 activity was detected visually by peripheral clear ring areas surrounding wells and distilled H₂O was used as negative control. Comparison of clear halos was measured in millimeters using UV transillumination (Gel Logic 200 Imagen System, Kodak, Rochester, NY, USA) and data were transformed to Phospholipase Units (PhU) in mm/μg.

Nicotinic acetylcholine receptor modulation assay

Electrophysiology was performed in *Xenopus laevis* oocytes expressing mouse fetal muscle ($\alpha 1\beta 1\gamma \delta$) or human neuronal ($\alpha 7$) nicotinic acetylcholine receptor (nACh) subtypes receptors. Two electrode voltage-clamp recordings were performed in a 30μl recording chamber, clamping the oocyte at -70 mV, and gravity perfused with ND96 solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl₂, and 5mM HEPES, pH 7.2). Acetylcholine

(ACh) gated currents were generated by 1sec pulses/min of 1-5μM ACh for $\alpha 1\beta 1\gamma \delta$ and 200μM ACh for $\alpha 7$ nAChR subtypes. *Crotalus m. nigrescens* venom and each fraction obtained by RP-HPLC were dissolved in 5μl (~200μg of toxin) of ND96. Venom fractions were allowed to equilibrate with the receptors in a static bath for 5min before pulsing with ACh. Subsequently, ACh pulses and ND96 perfusion were restarted to permit toxin dissociation from the receptors.

Vasculotoxicity

In vitro proteolysis of basement membrane

This was carried out as described previously (Escalante et al, 2011) using Matrigel™ (BD, United States) as basement membrane-like substrate. Matrigel was incubated with the venom in a venom:substrate ratio of 1:5 (w/w), for 15min, 1hr and 2hr at 37°C. Reactions were stopped by adding 5x SDS-PAGE sample loading buffer, then they were stored at -20°C until its utilization. Matrigel proteolysis was analyzed using a 4-12% (w/v) SDS-PAGE under reducing conditions.

Endothelial cells cytotoxicity and morphology changes

Median cytotoxic concentration (CC₅₀) induced by *C. m. nigrescens* venom was assessed by Lactate dehydrogenase (LDH) release into the media culture. Porcine aortic




Figure 4. Modulatory activity of *C. m. nigrescens* venom and its RP-HPLC fractions on nAChR expressed on *Xenopus laevis* oocytes. **A.** Percentage response was defined as current amplitude after application of *C. m. nigrescens* venom RP-HPLC fractions prior toxin application. **B.** Representative current trace elicited by one second pulse of ACh in muscle ($\alpha 1\beta 1\gamma \delta$) nAChR (black trace) and, in presence of venom fraction 1 (magenta trace). **C.** Current trace elicited by one second pulse of ACh in neuronal ($\alpha 7$) nAChR (black trace) and, in presence of fraction 19 in neuronal ($\alpha 7$) nAChR (red trace). **D.** *C. m. nigrescens* crude venom effect in neuronal ($\alpha 7$) nAChR, left trace represents control current elicited prior venom application. After 5 min equilibration period, ACh pulses resumed and ND96 solution was perfused over the oocyte between ACh pulses, with a loss of membrane potential.

endothelial (PAE) cells were kindly donated by Dr. Manuel Miranda-Arango (University of Texas at El Paso). The cells were grown in 5% (v/v) CO₂ at 37°C to 80% confluence and were treated with *C. m. nigrescens* venom (1-20µg/ml) diluted in F-12 media supplemented with 10% (v/v) fetal bovine serum and 100µg/ml gentamicin for 24hr. After venom incubation, media culture was collected and LDH release was measured by CytoTox 96® Non-Radioactive Cytotoxicity Assay (Promega, United States) according to the instructions of the manufacturer. The maximum LDH release was established by cell lysis using a Sonic Dismembrator 60 (Fisher Scientific, United States). Images of cell morphology were taken in an Axio Vert.A1 Microscope (Carl Zeiss, Germany).

Statistical analysis

For experiments with multiple repeated data sets, means and standard errors were calculated and ANOVA test ($P < 0.05$) was applied.

RESULTS

Hemotoxicity

In hemolysis assays we found that *C. m. nigrescens* venom produces Oxy-Hb release and Met-Hb formation (Figure 1). Oxy-Hb showed a median hemolytic concentration of $11.14 \pm 3.24 \mu\text{g/ml}$ and generated a full hemolysis at $40 \mu\text{g/ml}$ of *C. m. nigrescens* venom; Met-Hb showed a 50% formation at a $36.26 \pm 13.30 \mu\text{g/ml}$ of venom concentration

and generated a maximum formation at $320 \mu\text{g/ml}$ of *C. m. nigrescens* venom.

In order to investigate cell damage induced by *C. m. nigrescens* venom concentration we measured cell viability of human leukocyte cultures exposed in different concentrations of venom (0-10µg/ml) and different time intervals (2-12hr). Our results indicated that leukocyte cytotoxicity increased in time- and concentration-dependent manner. In the first 2hr of venom incubation, the cytotoxicity had the biggest increase with a 40% (Figure 2A) this result can be confirmed with CC_{50} values, where at 2hr was of $3.93 \pm 0.15 \mu\text{g/ml}$ and decreasing through time until CC_{50} decreased to $1.42 \pm 0.16 \mu\text{g/ml}$ in 12hr incubation (Figure 2B).

α -Neurotoxicity

A total of 19 fractions were collected from *C. m. nigrescens* venom by RP-HPLC, as are shown in Figure 3. Where the fractions 8, 9, 10, 11, 17, and 18 show specific phospholipase activity, and all displayed a higher activity than the *C. m. nigrescens* crude venom. Fractions 9, 10 and 11 had the highest phospholipase activity with 486.49, 237.30 and 738 PhU respectively. Followed by the fractions 8, 17, and 18 with 71.91, 21.80 and 25.92 PhU respectively. *C. m. nigrescens* venom had a phospholipase activity of 14.75 PhU.

The nAChR subtypes tested in this work ($\alpha 1\beta 1\gamma\delta$ and $\alpha 7$) presented a minimal or null modulation by *C. m. nigrescens*




Figure 5. Matrigel hydrolysis by *C. m. nigrescens* venom. **A.** SDS-PAGE of Matrigel incubated either with 0.9% (w/v) saline as control (CTL) or with venom (CMNv) during 0min, 15min, 1hr and 2hr. Lane 9 shows 50µg of crude CMNv. **B.** Densitogram correspondent to Figure 5A. Incubation times are represented in lines: control (black), 15min (red), 1hr (blue), and 2hr (green). Matrigel proteins are indicated by arrows: perlecan (1), laminin $\alpha 1$ (2), laminin $\beta 1$, laminin $\gamma 1$, type IV collagen (3) and nidogens (4, 5, and 6).

venom and the RP-HPLC fractions (Figure 4A). In $\alpha 1\beta 1\gamma\delta$ nAChR, the fractions 1, 2, and 14 generated less than 1% activity increase of the current (Figure 4B), and the fractions 6 and 9 generated a 3.33% receptor current decrease. In $\alpha 7$ nAChR, the fractions 3, 9, 13 and 19 (Figure 4C) generated 3.16% receptor current increase, and the fractions 4, 6, 11, and 16 generated 2.37% receptor current decrease. The *C. m. nigrescens* venom did not generate a modulation in both nAChR subtypes currents, but the resting potential of the oocyte after ACh pulses was not reconstituted as control after venom incubation (Figure 4D).




Figure 6. *C. m. nigrescens* venom cytotoxicity on PAE cells. PAE cells were treated with different concentrations of *C. m. nigrescens* venom (0-5 µg/ml) during 24hr. Data are represented as the mean of three independent experiments with their respective standard error.

Vasculotoxicity

When Matrigel was exposed to *C. m. nigrescens* venom, we observed degradation of many basement membrane constituents, depending of exposure time (Figure 5, A and B). The molecular masses of the different hydrolysates were calculated directly from the SDS-PAGE through R_f (Retardation factor). As control, six bands were observed in intact Matrigel: perlecan (>1,000 kDa); laminin $\alpha 1$ (420 kDa); laminin $\beta 1$, laminin $\gamma 1$, and type IV collagen (190 kDa); and nidogen (130, 110, and 90 kDa). 15min and 1hr incubation with *C. m. nigrescens* venom resulted in degradation of perlecan, all nidogen bands, and partial degradation of laminin $\alpha 1$, producing hydrolysis products of 370, 144, 100, 61, 54, 49, 40 and 21 kDa. After 2hr incubation all proteins of Matrigel were proteolyzed, except the collagen IV (190 kDa) band, which was partially degraded, producing hydrolysis products of 320, 160, 144, 132, 110, 82, 68, 56, 42, 36, and 21 kDa.

Crotalus m. nigrescens venom produced biphasic cytotoxicity in PAE cells: The first phase CC_{50} was $0.79 \pm 0.29 \mu\text{g/mL}$ (Figure 6) and is related to morphological changes in PAE cells as is shown in Figure 7B, where we observed cell detachment from substrate, starting from $0.1 \mu\text{g/ml}$ and cell thickness is reduced, becoming thinner than the control (Figure 7D and E); the second phase CC_{50} was $3.42 \pm 0.06 \mu\text{g/ml}$ (Figure 6), which was

morphologically observed as an aggregation of detached cells and cellular debris (Figure 7F, G and H).

DISCUSSION

Hemotoxicity

Hemolysis produced by *C. m. nigrescens* venom is already described (Macias-Rodríguez et al, 2014); even when the experimental procedures were different, the *C. m. nigrescens* venom had a concentration-dependent effect on erythrocytes, as we observe in our experiments. In this work, we demonstrate that the *C. m. nigrescens* venom generate a complex hemolytic process in a concentration-dependent manner, releasing Oxy-Hb release at low venoms concentrations and increasing Met-Hb when venom concentration is increased. There is not enough information about the mechanism of hemolytic process of this venom, but Oxy-Hb could be generated by destabilization and degradation of erythrocyte membrane via PLA2 (Doley et al, 2010), resulting in hemoglobin release. As a consequence, free hemoglobin can react with LAAO-produced H_2O_2 , resulting in oxidation of heme group and generating Met-Hb (Sunitha et al, 2015).

Hemolytic process during envenomation can cause oxygen transportation problems. Methemoglobin, a highly pro-oxidant molecule, can induce serious health problems in bite victims in the form of proinflammatory effect and oxidative stress. This can lead to long-term complications, such as thrombocytopenia, renal abnormalities and persistent local tissue degradation (Sunitha et al, 2015).

The leukocyte cytotoxicity could be a consequence of the disruption of the cellular membrane catalyzed by PLA2, as we discussed previously in the hemolytic process. Comparing with other cell types (erythrocytes and PAE cells) used to evaluate the effect of *C. m. nigrescens* venom, leukocytes had a higher susceptibility to the venom (lower CC_{50} of $1.42 \pm 0.16 \mu\text{g/ml}$) than in erythrocytes (CC_{50} of $11.14 \pm 3.24 \mu\text{g/ml}$) and PAE cells (CC_{50} of $3.42 \pm 0.06 \mu\text{g/ml}$). Also, leukocytes showed damage starting just after 2hr incubation, whereas in the erythrocytes and PAE cells no damage was observed. Even when the susceptibility is higher, none of the tested time intervals reached 100% of cytotoxicity. This phenomenon could be due to the activity of some isoforms of PLA2 induce adhesion to the endothelium instead of cytotoxicity (Zambelli et al, 2008).

α -Neurotoxicity

Crotalus m. nigrescens venom and its fractions did not modulate the current activity on fetal and $\alpha 7$ nAChR tested in this work. Even the fractions with phospholipase activity that in other species are found to produce α -neurotoxicity (such as crotoxin in *Crotalus durissus terrificus* venom (Sampaio et al, 2010), thus being similar to PLA2 RP-HPLC fractions of *C. m. nigrescens* venom) had no neurotoxic effect in nAChR.

C. m. nigrescens venom altered the resting potential of the oocytes via plasmatic membrane destabilization. This destabilization could be mediated by membrane peroxidation

due to L-amino acid oxidase H_2O_2 production (Torii et al, 1997) leading to the cell death process (necrosis or apoptosis) (Suhr and Kim, 1996). Also, PLA2 could contribute to the phenomena via phospholipid degradation (Doley et al, 2010).

Vasculotoxicity

Basement membrane-like components could be hydrolyzed by the catalytic activity of P-I and P-III SVMP. In other snake venoms, it was demonstrated that P-I SVMP has affinity for all the basement membrane previously listed (Escalante et al, 2011). On the other hand, P-III SVMP has an affinity for laminin $\beta 1$ and laminin $\gamma 1$ at longer periods of time than P-I SVMP (Escalante et al, 2006). The products of degradation generated by *C. m. nigrescens* venom could correspond to laminin $\gamma 1$ fragments (144 and 56 kDa bands), laminin $\alpha 1$ fragments (56 kDa band), and nidogen fragments (100, 56, 54, 42 and 41 kDa bands). The difference in degradation time of the components of basement membrane suggests that the hemorrhage formation could be generated in two steps: the first step of hemorrhage formation could be led by early degradation of perlecan and nidogens, forming dissociation of the basement membrane complex conformed by laminin and type IV collagen networks (Gutiérrez et al, 2005); the second step could be generated as a consequence of laminin and type IV collagen degradation, which could be related to endothelial cell detachment (Tanjoni et al, 2005).

Biphasic cytotoxic behavior of PAE cells, in contrast to morphology, denotes that the cytotoxicity could occur in two steps. The first step appears at low concentration of *C. m. nigrescens* venom (0.05-3.0 μ g/ml), where cytotoxicity is minimal but cell detachment occurs. This observation may have been due to SVMP and disintegrins which block the interaction of integrins with basement membrane components (Wu et al, 2003; Suntravat et al, 2015), or in this case, the interaction with the substrate. Also, in this interaction SVMP or disintegrin with integrins cause the cytoskeleton destabilization,

generating the cell thickness observed in Figure 7D and E (see Gutiérrez et al, 2005). The second step occurs at higher *C. m. nigrescens* venom concentrations (>3.0 μ g/ml) generating cell death; there are two possible scenarios for this: i) oxidative stress via LAAO H_2O_2 production (Samel et al, 2006), or ii) cellular membrane hydrolysis by PLA2 (Sampaio et al, 2010). Furthermore, other studies suggest that under these conditions the P-I SVMP could produce apoptosis anoikis (Tanjoni et al, 2005; Baldo et al, 2008).

CONCLUSIONS

C. m. nigrescens venom produced both Met-Hb formation and leukocyte cytotoxicity in a concentration-dependent manner. This venom and its HPLC-fractions did not generate α -neurotoxicity in muscle ($\alpha 1\beta 1\gamma \delta$) and neuronal ($\alpha 7$) nAChR subtypes. Finally, this venom hydrolyzed basement membrane-like substrate with a higher preference for nidogens and perlecan, and triggered PAE cells cytotoxicity.

ACKNOWLEDGEMENTS

This project was supported by Programa Integral de Fortalecimiento Institucional, SEP (PIFI) for Academic Groups, Universidad Autónoma de Ciudad Juárez. Electrophysiology experiments were partially supported by Consejo Nacional de Ciencia y Tecnología (CONACYT), Grant 153915. Also we want to acknowledge to Universidad Autónoma de Querétaro Herpetary (INE/CITES/DGVS-CR-IN-0619-QRO-00) for venom donation, Dr Edna Rico from Toxicology Lab of UACJ for donate the Matrigel used, Dr Manuel Miranda-Arango from Border Biomedical Research Center at UTEP for cell PAE cells donation and Dr Angel Diaz for his criticism in this paper.

COMPETING INTERESTS

None declared.




Figure 7. Morphological changes induced by *C. m. nigrescens* venom on PAE cells. Optic microscopy of PAE cells treated with different concentrations of *C. m. nigrescens* venom: a = 0.0 μ g/ml, b = 0.1 μ g/ml, c = 0.5 μ g/ml, d = 1.0 μ g/ml, e = 2.0 μ g/ml, f = 3.0 μ g/ml, g = 4.0 μ g/ml, and h = 5.0 μ g/ml. Symbols: death cells (black arrow), cellular thickness decrease (red arrow), cellular aggregation (star).

ABBREVIATIONS

Ach; Acetylcholine
 ACN; Acetonitrile
 CC₅₀; Median cytotoxic concentration
 DMEM; Dulbecco's Modified Eagle Medium
 LAAO; L-amino acid oxidase
 LDH; Lactate dehydrogenase
 nAChR; Nicotinic acetylcholine receptor
 PAE; Porcine aortic endothelial cells
 PhU; Phospholipase units
 PLA2; Phospholipase A2
 RBC; Red blood cells
 RP-HPLC; Reverse phase-High Performance Liquid Chromatography
 SVMP; Snake venom metalloproteinases
 TFA; Trifluoroacetic acid

REFERENCES

- Bon C, Changuex JP, Jeng TW and Fraenkel-Conrat H. 1979. Postsynaptic effects of crotoxin and of its isolated subunits. *Eur J Biochem*, 99, 471-481.
- Braga MDM, Costa Martins AM, Amora DN, et al. 2008. Purification and biological effects of L-amino acid oxidase isolated from *Bothrops insularis* venom. *Toxicon*, 51, 199-207.
- Das D, Urs N, Hiremath V, Vishwanath BS and Doley R. 2013. Biochemical and biological characterization of *Naja kaouthia* venom from North-East India and its neutralization by polyvalent antivenom. *J Venom Res*, 4, 31-38
- Doley R, Zhou X and Kini M. 2010. Snake venom phospholipase A2 enzymes. In: Mackessy SP (Ed) *Handbook of venoms and toxins of reptiles*, CRC Press, Florida, United States of America, 1st edition, pp 174-205.
- Escalante T, Shannon JD, Moura da Silva AM, Gutierrez JM and Fox JW. 2006. Novel insights into capillary vessel basement membrane damage by snake venom hemorrhagic metalloproteinases: a biochemical and immunocyto-chemical study. *Arch Biochem Biophys*, 455, 144-153.
- Escalante T, Ortiz N, Rucavado A et al. 2011. Role of Collagens and perlecan in microvascular stability: Exploring the mechanism of capillary vessel damage by snake venom metalloproteinases. *PLoS ONE*, 6, e28017.
- Gutiérrez JM, Rucavado A, Escalante T and Díaz C. 2005. Hemorrhage induced by snake venom metalloproteinases: biochemical and biophysical mechanisms involved in microvessel damage. *Toxicon*, 45, 997-1011.
- Hardy DL, Jeter M and Corrigan Jr. JJ. 1982. Envenomation by the northern blacktail rattlesnake (*Crotalus molossus molossus*): report of two cases and the in vitro effects of the venom on fibrinolysis and platelet aggregation. *Toxicon*, 20, 487-493.
- Kattah LS, Santoro MM, Diniz CR and De Lima ME. 2000. Crotoxin, the major toxin from the rattlesnake *Crotalus durissus terrificus*, inhibits 3H-choline uptake in guinea pig ileum. *Braz J Med Biol Res*, 33, 1093-1097.
- Lemos-Espinal JA and Smith HM. 2009. *Anfibios y reptiles del estado de Chihuahua, México*, Universidad Nacional Autónoma de México and Comisión Nacional para el Conocimiento y uso de la Biodiversidad, México City, México. 1st edition, 222-223
- Lowry OH, Rosebrough NJ, Farr AL and Randall RJ. 1951. Protein measurement with the Folin phenol reagent. *J Biol Chem*, 193, 265-275.
- Macias-Rodríguez EF, Martínez-Martínez A, Gatica-Colima A, Bojórquez-Rangel G and Plenge-Tellechea LF. 2014. Análisis comparativo de la actividad hemolítica entre las subespecies *Crotalus molossus molossus* y *Crotalus molossus nigrescens*. *Revista Bio Ciencias*, 2, 302-312.
- Mackessy SP. 2000. The field of reptile toxicology: snakes, lizards, and their venoms. In: Mackessy SP (Ed) *Handbook of venoms and toxins of reptiles*, CRC Press, Florida, United States of America, 1st edition, pp 3-23.
- Meléndez-Martínez DL, Macias-Rodríguez E, Vargas-Caraveo A et al. 2014. Capillary damage in the area postrema by venom of the black-tailed rattlesnake (*Crotalus molossus molossus*). *J Venom Res*, 5, 1-5.
- Ramírez GA, Fletcher Jr. PL and Possani DL. 1990. Characterization of the venom from *Crotalus molossus nigrescens* Gloyd (Black tail rattlesnake): Isolation of two proteases. *Toxicon*, 28, 285-297.
- Samel M, Vija H, Ronnholm G et al. 2006. Isolation and characterization of an apoptotic and platelet aggregation inhibiting L-amino acid oxidase from *Vipera berus berus* (common viper) venom. *Biochim Biophys Acta*, 1764, 707-714.
- Sampaio SC, Hyslop S, Fontes MR et al. 2010. Crotoxin: novel activities for a classic beta-neurotoxin. *Toxicon*, 55, 1045-1060.
- Suhr SM and Kim DS. 1996. Identification of the snake venom substance that induces apoptosis. *Biochem Biophys Res Commun*, 224, 134-139.
- Sunitha K, Hemshekhar M, Thushara RM et al. 2015. Inflammation and oxidative stress in viper bite: An insight within and beyond. *Toxicon*, 98, 87-98.
- Suntravat M, Barret HS, Jurica CA et al. 2015. Recombinant disintegrin (r-Cam-dis) from *Crotalus adamanteus* inhibits adhesion of human pancreatic cancer cell lines to laminin-1 and vitronectin. *J Venom Res*, 6, 1-10.
- Tanjoni I, Weinlich R, Della-Casa MS et al. 2005. Jararhagin, a snake venom metalloproteinase, induces a specialized form of apoptosis (anoikis) selective to endothelial cells. *Apoptosis*, 10, 851-861.
- Torii S, Naito M and Tsuruo T. 1997. Apoxin I, a novel apoptosis-inducing factor with L-amino acid oxidase activity purified from western diamondback rattlesnake venom. *J Bio Chem*, 272, 9539-9542.
- Valdes JJ, Thompson RG, Wolff VL et al. 1989. Inhibition of calcium channel dihydropyridine receptor binding by purified Mojave toxin. *Neurotoxicol Teratol*, 11, 129-133.
- Verma RS and Babu A. 1995. *Human Chromosomes Principles and Techniques*, McGraw-Hill Inc, New York, United States of America, 2nd edition.
- Wu WB, Peng HC and Huang TF. 2003. Disintegrin causes proteolysis of β -catenin and apoptosis of endothelial cells: Involvement of cell-cell and cell-ECM interactions in regulating cell viability. *Exp Cell Res*, 286, 115-127.
- Yarema MC and Curry SC. 2005. Envenomation by the northern blacktail rattlesnake (*Crotalus molossus molossus*) case report. *Pediatr Emerg Care*, 21, 40-42.
- Zambelli VO, Sampaio SC, Sudo-Hayashi LS et al. 2008. Crotoxin alters lymphocyte distribution in rats: Involvement of adhesion molecules and lipoxigenase-derived mediators. *Toxicon*, 51, 1357-1367.